

Modelling of Alfvén modes properties in TJ-II plasmas

A. Rakha¹, M. J. Mantsinen^{1,2}, A. V. Melnikov^{3,4}, S. E. Sharapov⁵, D. A. Spong⁶,
A. Gutiérrez-Milla¹, A. López-Fraguas⁷, F. Castejón⁷, J. L. de Pablos⁷, and X. Sáez¹

¹Barcelona Supercomputing Center, Spain

²ICREA, Barcelona, Spain

³National Research Center 'Kurchatov Institute', 123182, Moscow, Russia

⁴National Research Nuclear University MEPhI, 115409, Moscow, Russia

⁵CCFE, Culham Science Centre, OX14 3DB, UK

⁶Oak Ridge National Laboratory, Oak Ridge, TN-37831, USA

⁷Fusion National Laboratory, CIEMAT, 28040, Madrid, Spain

Alfvén modes are a low-frequency electromagnetic class of instabilities, which degrade the fast ion confinement and result in poor performance. In burning plasma devices, i.e. ITER and beyond, it is expected that energetic alpha particles will heat the bulk plasma; therefore good confinement and the interaction of energetic particles with background plasmas are of the most important interest in fusion research. The study of Alfvén Eigenmodes (AE) in tokamak plasmas [1] and in 3D non-axisymmetric devices like TJ-II [2,3] demands detailed investigations and modelling of these structures for the optimization of fusion devices. The study of the properties of AE modes and their effect on fast ion confinement is deemed necessary for the enhanced performance of next step burning plasmas [4]. In this work, we take advantage of TJ-II flexibility and present modelling of the observed behavior of Alfvénic modes in a configuration scan of TJ-II, shown in Figure 1 from [5], and compare these results with the experimental ones for the different properties of Alfvén Eigenmodes in TJ-II plasmas. Simulations of selected instabilities [5] are performed by using STELLGAP [6] and AE3D [7] codes for the three-dimensional flexible Helic TJ-II. Simulation results for the properties of observed AE modes i.e. frequency and radial location are consistent with the experimental observations. Additionally, in our simulations, the mode with same poloidal and toroidal mode numbers appears with the small variations in on-axis iota values. In next step, we will investigate the transition of AE modes between chirping and steady at different iota values.




Figure 1: Experimental observations of Alfvén Eigenmodes (AE) in TJ-II Stellarator.

References

- [1] Fasoli A. et al 1995 Nucl. Fusion 35 1485
- [2] Ascasibar E. et al. 2002 Plasma Phys. and Contr. Fusion 44 B307
- [3] Jiménez-Gómez R. et al. , Nucl. Fusion 51,033001 (2011)
- [4] Gorelenkov N.N. et al 2003 Nucl. Fusion 43 594
- [5] Melnikov A.V., et. al. 2016 Nucl. Fusion 56 076001
- [6] Spong D. A., Sanchez R. and Weller A. 2003 Phys. Plasmas 10 3217
- [7] Spong D. A., Azevedo D' and Todo Y. 2010 Phys. Plasmas 17 022106