

Scrape-off layer simulations in a Double Null magnetic configuration

C. F. Beadle, P. Ricci, P. Paruta, F. Riva, C. Wersal

Swiss Plasma Centre, EPFL

Motivation for Double Null (DN) configuration

Simulation of Double Null SOL

Main features of the Double Null SOL



ÉCOLE POLYTECHNIQUE
FÉDÉRALE DE LAUSANNE


**SWISS PLASMA
CENTER**

The Tokamak SOL

Cross-field transport driven by turbulence

Rapid parallel outflow

$L_{\text{SOL}} \sim 1\text{cm}-1\text{mm}$



Heat exhaust

Impurity control


Plasma confinement

Plasma fuelling

Regulating neutral density

Ashes removal ($\text{He} < 10\%$)

DN: better power handling?



Share power evenly
between the two outer legs

Possibility of two radiation
fronts in detached operation

DN: HFS ICRF launchers?

Quiescent HFS


Antenna and launchers here?

Calls for
theoretical
investigation!

plasma without damage

➤ Efficient coupling

GBS: our simulation code



GBS: our simulation code


2 fluid Drift Reduced Braginskii Equations
evolving $n, \phi, T_e, v_{\parallel e}, v_{\parallel i}$

Flux driven: no separation between equilibrium
and fluctuations

First principles boundary conditions


Finite differences, 4th order RK timestepping

From GBS past work in limited configuration to
Double Null




**Solution:
geometrical
coordinates**

Double null implemented in geometrical coordinates



Flux function based on 3 current carrying wires



Up-down and poloidally left-right symmetric

Toroidicity effects are included


$q \simeq 3.5$ at inner boundary

At LCFS $q \rightarrow \infty$ and $L_c \rightarrow \infty$


Strong magnetic shear close to X points

B field include $1/R$ dependence





First results – density evolution in poloidal plane





First results – electrical potential evolution in the poloidal plane



Complex Equilibrium and Fluctuation Pattern




Upper X: Circulation around X point



ExB drifts important effect on target heat flux
Corresponds to comparison between
experiment and phenomenological simulations

Lower X: Symmetric divertor legs with low turbulence






Dominant flow along outer leg

Strong shear across leg



Low turbulence on leg

LFS: minimum pressure spatial decay length at midplane


$$p_e = p_{e0} e^{(-r/L_{pr})} = p_{e0} e^{-(\psi - \psi_0)/L_{p\psi}}$$




LFS: narrow feature at midplane



$\ln(p_e/p_{e0})$




Single scale length



More than one scale length






Decay with ψ takes similar form



Single scale length

r/ρ_s

LFS: Narrow feature due to strong ExB flow shear



$$\omega_{E \times B} \sim \gamma_{BM}$$




ExB shear flow driven by high potential fluctuations

HFS: very little plasma outside LCFS

No interchange drive

Pressure at LCFS 10x lower than LFS

Very strong flow shear -> KH instability?



Concluding Remarks

- First simulations of plasma turbulence in the SOL of a double null configuration
- Leveraging upgrade to GBS simulation code
- Initial analysis shows a complex flow pattern
- Up-down and left-right asymmetry in divertor legs
- Highly turbulent LFS with narrow feature at midplane
- Quiescent, almost empty HFS

Questions?