




Retrospects and prospects of X-ray cluster surveys



Rosati, Norman, Borgani 2002

Alexis Finoguenov



X-ray cluster identification

Point source+by eye BCS, REFLEX, NORAS, MACS	Point source+optical survey CODEX, RASSinDES
Extended source+by eye RDCS, 400 sq.deg., SPT	Extended source+optical survey COSMOS, CDFS, AEGIS, SXDF, XMM-XXL

10^{-13}

1.1368684E-13

2

-2.2737368E-13

10¹⁴





4


-4.5474735E-13

8


-9.084947E-13

Sensitivity of RASS on BOSS footprint






CODEX and MCXC




RASS vs Redmapper mass estimates




$z: [0.3, 0.4]$

X-Ray M500




Optical M500



z: [0.3, 0.4]




X-Ray M500



z: [0.4, 0.5]


z: [0.5, 0.6]

Optical M500



SDSS-IV surveys (2014 through 2020)


APOGEE-2 



Maps hundreds of thousands of individual stars in the Milky Way.


galaxy evolution & dark matter

MaNGA 



Maps 10,000 nearby galaxies.


**eBOSS,
TDSS,
SPIDERS** 



Maps the Universe of galaxies and quasars. Especially quasars.

dark energy & cosmology

Galaxy clusters: status ($1 \text{ yr}^{3/4}$)




Status 2016-03-29 (CODEX)

1942 candidates richness > 10


- 1307 $N_{spec} > 3$
- 754 $N_{spec} > 10$
- 427 $N_{spec} > 15$


Of them: 1600/1942 completed

- 1051 $N_{spec} > 3$
- 630 $N_{spec} > 10$
- 369 $N_{spec} > 15$



CFHT-CODEX 27-best sample








Cibirca+


W
L

m
o
d
e
|
10





LoCuSS, CLASH, CODEX





LoCuSS, CLASH, CODEX



AGN contamination line: $\langle L_{X\text{AGN}} \rangle * \text{HOD}$

Ebrero XLF(z) + Allevato HOD

